



VICTORIA CARDIOLOGY

ACCREDITED ECHOCARDIOGRAPHY LABORATORY



Unit 201, 1252 Lawrence Ave. East, North York, ON M3A 1C3

Tel: 416.609.1100 Fax: 416.609.1102

Dr. A. Janmohamed MBBS., FRCP(C) Dr. M. Zia MD, FRCP(C)

Dr. K. Merali MD, FASE Dr. R. Poblete MD, FRCP(C)

Dr. M. Sud MD, FRCP(C) Dr. S. Miller MD, FRCP(C)

Appointment: _____

PATIENT INFORMATION (Please label here if applicable)	
Patient Name _____	<input type="checkbox"/> F <input type="checkbox"/> M
OHIP# _____	Version Code _____
Home Address _____	
Phone# _____	DOB: _____

CARDIOLOGY CONSULTATION AND APPROPRIATE TESTING

- | | |
|--|---|
| <input type="checkbox"/> Cardiology Consultation | <input type="checkbox"/> ECG |
| <input type="checkbox"/> Echocardiogram <input type="checkbox"/> Contrast Echo | <input type="checkbox"/> 24/48/72 Hours Digital Holter Monitoring |
| <input type="checkbox"/> Stress Echo (Treadmill Exercise and Echo) | <input type="checkbox"/> 7/14 Days Digital Holter Monitoring |
| <input type="checkbox"/> Contrast Stress Echo | <input type="checkbox"/> Consult if Test is Abnormal |
| <input type="checkbox"/> Stress ECG Test (Treadmill Exercise) | <input type="checkbox"/> ***24hrs Ambulatory Blood Pressure Monitoring*** |

INDICATIONS FOR STRESS ECHO

- Typical or atypical chest pain of ischemic equivalent syndrome
- Possible ACS with non-diagnostic ECG changes and negative or borderline significant troponin levels
- History of Congestive Heart Failure
- Known LV systolic dysfunction of unclear etiology
- Significant ventricular arrhythmia
- Syncope of unclear etiology
- Borderline or high troponin levels in a setting other than ACS
- Significant cerebrovascular or peripheral atherosclerosis
- Re-evaluation (≥ 1 yr) in patients with significant cerebrovascular or peripheral atherosclerosis
- Equivocal or non-diagnostic results from other stress modalities
- Initial evaluation of patients at intermediate or high global CAD risk
- Periodic (≥ 2 yrs) re-evaluation of patients with intermediate or high global CAD risk
- New or worsening chest pain or ischemic equivalent.
- Post MI or ACS for risk stratification (within 3 months).
- Viability in patients with known significant LV dysfunction post re-vascularization
- Periodic (≥ 1 yr) re-evaluation of stable patients with known CAD (previous coronary angiography, CTA/EBCT, MI, ACS or abnormal stress imaging)
- For physiologic assessment and/or symptoms correlation in patients with moderate or severe Aortic Stenosis, Mitral Stenosis, Mitral Regurgitation, Aortic Regurgitation, Hypertrophic Cardiomyopathy
- Assessment of established or latent pulmonary hypertension
- Others:**

INDICATIONS FOR ECHOCARDIOGRAPHY

- Heart Murmurs
- Native Valvular Stenosis
- Native Valvular Regurgitation
- Known or Suspected Mitral Valve Prolapse
- Congenital or inherited Cardiac Structural Disease (including Bicuspid Aortic Valve, Marfan's Syndrome, Atrial Septal Defect, Ventricular Septal Defect, Ehler's Danlos Syndrome)
- Prosthetic Heart Valves
- Infective Endocarditis
- Pericardial Disease
- Cardiac Masses
- Interventional Procedures
- Pulmonary Diseases
- Chest Pain and Coronary Artery Disease
- Dyspnea, Edema and Cardiomyopathy
- Hypertension
- Thoracic Aortic Disease
- Neurologic or other Possible Embolic Events
- Arrhythmias Syncope and Palpitations
- Before Cardioversion
- Suspected Structural Heart Disease
- Others:**



Clinical Information: (Please fax previous report if available)

HIGH-RISK STUDIES WILL BE COMMUNICATED OVER THE PHONE

Referring Physician _____ Provider Billing #: _____

Verbal Fax: _____ Signature: _____

VICTORIA CARDIOLOGY

Unit 201, 1252 Lawrence Ave. East, North York, ON M3A 1C3
 Tel: 416.609.1100 Fax: 416.609.1102



CARDIAC TEST PREPARATION AND INFORMATION

CONSULTATION:

Please bring all your medications.

24 / 48 HOURS HOLTER MONITORING:

No creams/lotion on the chest. 24 and 48 hours monitoring is **NON-STOP** recording. Please be prepared for **NO SHOWERING or SWIMMING** during the next 24 or 48 hours.

72 HOURS AND 7-DAY / 14-DAY HOLTER MONITORING:

No cream / lotion on the chest. 72 hours and 7/14-day monitoring allows patients to take a shower with Monitor Removal. Patients will be taught how to put back electrodes after shower. Monitor hook-up takes 15 to 20 minutes.

ECHOCARDIOGRAM:

No preparation. Test takes 35-45 minutes.

STRESS ECHO / STRESS TEST:


Please wear loose clothing and running shoes. Ask your doctor if you should be stopping any of your medication. Do not apply body cream/lotion on the day of the test. Test takes 45-60 minutes.

*****24hrs Ambulatory Blood Pressure Monitoring***:**

Not insured by OHIP. \$70 charge applies.



HOW TO FIND US



Check our website for more info: www.victoriamedicalgroup.ca